

MALE 1: Eloctate, antihemophilic factor, recombinant, FC fusion protein step by step reconstitution video. Hi, I'm Jonathon. I know many of you have been treating your hemophilia A for a long time and I get that you're used to infusing a certain way. I'm sure your doctor or nurse showed you how to do this. However, I'm going to show you step by step how to reconstitute Eloctate. At the end of this video, you should feel ready to reconstitute with this process.

[00:00:31]

Talk to your doctor with any questions, and always make sure to refer to the instructions for use. The Eloctate reconstitution device is packaged together so everything you need is at your fingertips. Now let's get started. To begin, locate the arrows on both sides on the bottom of the kit and pinch them. Then lift up on the half circles and the kit will open. Inside your kit you'll find a vial with a powdered drug, a vial adapter in its package, a pre-filled diluent syringe, and a plunger rod.

[00:01:06]

Take out the Eloctate vial and remove the plastic cap. Clean the rubber stopper with an alcohol wipe. Let it dry. Remove the backing from the vial adapter package by peeling back the lid. Place the vial on a flat surface.

[00:01:32]

Hold the vial with one hand, then using the other hand place the adapter over the vial. The spike should be placed directly above the center of the rubber stopper. Next, push the adapter straight down until the spike punctures the center of the vial stopper and is fully inserted. Now lift the package cover away from the adapter and discard the cover. Locate the plunger rod and hold it at the circular disk.

[00:02:00]

Place the tip of the plunger rod into the end of the syringe. Turn it clockwise until it's securely attached. With one hand, hold the diluent syringe right under the cap. Make sure the cap is pointing up and you're holding the diluent syringe by the ridged part directly under the cap. Take your other hand, grasp the cap and bend it at a 90 degree angle until it snaps off.

[00:02:32]

After the cap comes off, you will see the glass tip of the syringe. Be sure the vial is sitting on a flat surface. Now inset the tip of the syringe into the adapter opening. Turn the syringe clockwise until it's securely attached. Slowly press down the plunger rod to inject all the diluent into the vial. The plunger rod may rise slightly after this process. Don't worry, this is normal.

[00:03:00]

Keep the syringe connected to the adapter and gently swirl the vial until everything is dissolved. The appearance of the solution should be clear to slightly opalescent and colorless. If you're using more than one vial, stop here and refer to the pooling instructions in the instructions for use. Make the sure the plunger rod is completely depressed and then turn the vial upside down. Slowly, pull out the plunger rod to draw up the solution into the syringe.

[00:03:32]

Gently unscrew the syringe from the vial adapter. Dispose of the vial with the adapter still attached. Now you're ready to go. And always keep in mind reconstituted Eloctate should be administered as soon as possible. If the product is not used within three hours after reconstitution, it must be discarded.

[00:04:00]

And don't forget, always talk to your doctor or nurse if you have any questions.

MALE 2: Indications and important safety information. Indications: Eloctate antihemophilic factor, recombinant FC Fusion protein is a recombinant DNA-derived antihemophilic factor indicated in adults and children with hemophilia A, congenital factor VIII deficiency for on demand treatment and control of bleeding episodes, perioperative management of bleeding and routine prophylaxis to reduce the frequency of bleeding episodes.

[00:04:31]

Eloctate is not indicted for the treatment of Von Willebrand disease. Important safety information. Do not use Eloctate if you have had an allergic reaction to it in the past. Tell your healthcare provider if you have or have had any medical problems, take any medicines including prescription and non-prescription medicines, supplements or herbal medicines, have any allergies, are breastfeeding, or pregnant or planning to become pregnant, or have been told you have inhibitors, antibodies to factor VIII.

[00:05:02]

Allergic reactions may occur with Eloctate. Call your healthcare provider or get emergency treatment right away if you have any of the following symptoms: Difficultly breathing, chest tightness, swelling of the face, rash, or hives. Your body can also make antibodies called inhibitors against Eloctate which may stop Eloctate from working properly. The most frequently occurring side effects of Eloctate are headache, rash, joint pain, muscle pain and general discomfort. These are not all the possible side effects of Eloctate.

[00:05:31]

Talk to your healthcare provider right away about any side effect that bothers you or that does not go away, and if bleeding is not controlled after using Eloctate. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call 1-800-FDA-1088. Please see full prescribing information.

[00:06:03]